



Rambler Ranger Broiler

PERFORMANCE OBJECTIVES

2018



An Aviagen Brand



Contents

01	<i>Key Management Points</i>
02	<i>Introduction</i>
03	Section g <i>As-Hatched Performance</i>
04	Section g <i>Male Performance</i>
05	Section g <i>Female Performance</i>
07	Section lb <i>As-Hatched Performance</i>
08	Section lb <i>Male Performance</i>
09	Section lb <i>Female Performance</i>
11	<i>Carcass Yield</i>

Key Management Points

The Rambler Ranger™ broiler is a non-feather sexable, slower-growing colored broiler. Ideally suited for organic and free-range production, the Rambler Ranger is a very slow growing bird which offers good meat yield and exceptional health and welfare characteristics. The following points are important for optimizing performance of the Rambler Ranger broiler:

- Maximize chick quality by good management of hatching, storage and transport conditions.
- Design the brooding set-up to ensure easy access to water and feed at placement, and to ease the transition between supplementary systems and the automated feeders and drinkers at 4-5 days. Feed a highly digestible and nutritionally balanced Starter diet.
- Rambler Ranger broiler chicks might be slightly smaller as egg size of the Ranger female is lower than that of standard Aviagen® parent stock females. Early environmental temperatures might have to be adjusted to account for this (+ 1°C / 2°F higher).
- Keep chicks in their thermal comfort zone by monitoring chick behavior, but beware of low relative humidities (less than 50% RH). Establish a minimum ventilation program from day one.
- Monitor crop fill, feeding and drinking behavior and 7-day live weight to allow continuous improvement of the brooding set-up.
- Maintain high standards of biosecurity and cleanliness to keep disease to a minimum.



Introduction

This booklet contains the performance objectives for the Rambler Ranger broiler and is to be used in conjunction with the **Broiler Management Handbook** supplied by Aviagen.

Performance

The performance objectives given in this booklet will be achieved under good management and environmental conditions. The achieved daily gain depends on feed program and management applied, and climatic conditions. Different growing programs and market demands may require alternative feed formulations; the Aviagen nutrition team can provide advice to suit individual growing programs.

The objectives are presented in two sections to reflect the global nature of the publication.

Section 1 *g* contains the performance data in metric measurement and

Section 2 *lb* contains imperial measurements.

In the tables values are rounded, this may result in small inaccuracies when using the objectives to calculate other performance statistics.

Yield will vary between processing plants depending on type of equipment used (e.g. carcass chilling technology, automated versus manual deboning) and the exact portion being produced.

For further information on the management of Rambler Ranger broiler stock, please contact your local Aviagen Representative.



As-Hatched Performance

Day	Weight (g) ¹	Daily gain (g)	Av. daily gain / week (g)	Daily intake (g)	Cum. intake (g) ²	FCR ³
0	29					
1	35	6			8	0.224
2	43	8		10	18	0.405
3	53	10		12	29	0.550
4	64	11		14	43	0.667
5	77	12		16	59	0.762
6	91	14		18	76	0.840
7	106	15	10.96	20	96	0.905
8	122	17		22	117	0.960
9	140	18		24	141	1.007
10	160	19		26	167	1.048
11	181	21		28	196	1.084
12	203	22		31	226	1.116
13	227	24		33	260	1.145
14	252	25	20.94	36	296	1.172
15	279	27		38	334	1.197
16	307	28		41	375	1.221
17	337	30		44	419	1.243
18	368	31		47	466	1.265
19	400	32		49	515	1.286
20	434	34		52	567	1.307
21	469	35	30.94	55	622	1.327
22	505	36		58	680	1.347
23	542	37		61	741	1.367
24	580	38		63	804	1.386
25	619	39		66	870	1.405
26	659	40		69	939	1.424
27	700	41		71	1010	1.443
28	741	41	38.88	74	1084	1.462
29	783	42		76	1160	1.481
30	826	43		79	1238	1.500
31	869	43		81	1319	1.518
32	913	44		83	1402	1.536
33	957	44		85	1488	1.555
34	1001	44		87	1575	1.573
35	1046	45	43.56	89	1664	1.591
36	1091	45		91	1755	1.609
37	1136	45		93	1848	1.627
38	1181	45		94	1942	1.644
39	1226	45		96	2037	1.662
40	1271	45		97	2134	1.680
41	1316	45		98	2233	1.697
42	1360	45	44.92	99	2332	1.714
43	1405	45		100	2433	1.731
44	1449	44		101	2534	1.748
45	1493	44		102	2636	1.765
46	1537	44		103	2739	1.782
47	1580	43		104	2843	1.799
48	1623	43		104	2947	1.816
49	1666	42	43.59	105	3051	1.832
50	1708	42		105	3156	1.848
51	1749	42		105	3261	1.865
52	1790	41		105	3366	1.881
53	1831	40		105	3472	1.897
54	1870	40		105	3577	1.912
55	1910	39		105	3682	1.928
56	1949	39	40.42	105	3787	1.944
57	1987	38		105	3892	1.959
58	2024	38		105	3997	1.974
59	2061	37		104	4101	1.990
60	2097	36		104	4205	2.005
61	2133	36		103	4308	2.020
62	2168	35		103	4411	2.035
63	2202	34	36.24	102	4513	2.049
64	2236	34		102	4614	2.064
65	2269	33		101	4715	2.078
66	2301	32		100	4815	2.093
67	2333	32		99	4915	2.107
68	2364	31		99	5013	2.121
69	2394	30		98	5111	2.135
70	2424	30	31.68	97	5208	2.149

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.

NOTE: In the table values are rounded, this may result in small inaccuracies when using the objectives to calculate other performance statistics.



Rambler Ranger Broiler Performance Objectives



Male Performance

Day	Weight (g) ¹	Daily gain (g)	Av. daily gain / week (g)	Daily intake (g)	Cum. intake (g) ²	FCR ³
0	29					
1	35	6			7	0.205
2	43	8		9	16	0.375
3	53	10		11	27	0.516
4	64	11		13	41	0.633
5	76	12		15	56	0.731
6	90	14		18	73	0.814
7	105	15	10.92	20	93	0.883
8	122	17		22	115	0.943
9	141	18		24	140	0.994
10	160	20		27	167	1.039
11	182	21		29	196	1.078
12	205	23		32	228	1.113
13	230	25		35	263	1.145
14	256	26	21.51	38	301	1.174
15	284	28		40	341	1.200
16	314	30		43	384	1.225
17	345	31		46	430	1.248
18	378	33		49	480	1.271
19	412	34		52	532	1.292
20	447	36		55	587	1.312
21	484	37	32.63	58	645	1.332
22	523	38		61	707	1.351
23	563	40		64	771	1.370
24	603	41		67	838	1.389
25	646	42		70	908	1.407
26	689	43		73	981	1.425
27	733	44		76	1057	1.443
28	778	45	41.91	79	1136	1.461
29	824	46		82	1218	1.478
30	870	47		84	1302	1.496
31	918	47		87	1389	1.513
32	966	48		89	1478	1.530
33	1014	49		92	1569	1.547
34	1063	49		94	1663	1.564
35	1113	49	47.81	96	1759	1.581
36	1162	50		98	1857	1.598
37	1212	50		100	1957	1.615
38	1262	50		102	2059	1.631
39	1312	50		104	2163	1.648
40	1362	50		105	2268	1.665
41	1413	50		107	2375	1.681
42	1463	50	50.03	108	2483	1.697
43	1513	50		109	2592	1.713
44	1562	50		110	2702	1.730
45	1612	50		111	2814	1.746
46	1661	49		112	2926	1.762
47	1710	49		113	3039	1.777
48	1759	48		114	3153	1.793
49	1807	48	49.12	114	3268	1.809
50	1854	48		115	3383	1.824
51	1901	47		115	3498	1.840
52	1948	47		116	3613	1.855
53	1994	46		116	3729	1.870
54	2039	45		116	3845	1.885
55	2084	45		116	3961	1.901
56	2128	44	45.97	116	4077	1.915
57	2172	44		116	4192	1.930
58	2215	43		115	4308	1.945
59	2257	42		115	4423	1.960
60	2299	42		115	4538	1.974
61	2339	41		114	4652	1.989
62	2379	40		114	4766	2.003
63	2419	39	41.50	113	4879	2.017
64	2458	39		113	4992	2.031
65	2495	38		112	5104	2.045
66	2533	37		111	5215	2.059
67	2569	36		111	5326	2.073
68	2605	36		110	5436	2.087
69	2640	35		109	5544	2.100
70	2674	34	36.47	108	5653	2.114

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.

NOTE: In the table values are rounded, this may result in small inaccuracies when using the objectives to calculate other performance statistics.



Female Performance

Day	Weight (g) ¹	Daily gain (g)	Av. daily gain / week (g)	Daily intake (g)	Cum. intake (g) ²	FCR ³
0	29					
1	35	6			9	0.248
2	44	8		10	19	0.439
3	54	10		12	32	0.587
4	65	11		14	46	0.703
5	77	12		16	61	0.795
6	91	14		17	79	0.868
7	106	15	10.95	19	98	0.928
8	122	16		21	119	0.977
9	139	18		23	142	1.019
10	158	19		25	167	1.056
11	179	20		27	195	1.088
12	200	22		29	224	1.117
13	224	23		32	256	1.144
14	248	24	20.32	34	290	1.169
15	274	26		37	326	1.193
16	301	27		39	365	1.216
17	329	28		42	407	1.238
18	358	29		44	451	1.260
19	389	30		47	498	1.281
20	420	32		49	547	1.302
21	453	33	29.25	52	599	1.323
22	486	34		54	653	1.343
23	521	34		57	710	1.364
24	556	35		59	769	1.384
25	592	36		62	831	1.404
26	629	37		64	895	1.424
27	666	37		66	962	1.444
28	704	38	35.89	69	1031	1.464
29	742	38		71	1101	1.484
30	781	39		73	1174	1.504
31	820	39		75	1249	1.523
32	860	39		77	1326	1.543
33	900	40		79	1405	1.562
34	939	40		80	1485	1.581
35	980	40	39.37	82	1567	1.600
36	1020	40		84	1651	1.619
37	1060	40		85	1736	1.638
38	1100	40		86	1822	1.657
39	1140	40		87	1910	1.675
40	1180	40		89	1998	1.694
41	1219	40		90	2088	1.712
42	1259	39	39.87	90	2178	1.731
43	1298	39		91	2269	1.749
44	1337	39		92	2361	1.767
45	1375	39		93	2454	1.784
46	1413	38		93	2547	1.802
47	1451	38		94	2641	1.820
48	1488	37		94	2735	1.837
49	1525	37	38.09	94	2829	1.855
50	1562	36		94	2923	1.872
51	1598	36		94	3018	1.889
52	1633	35		94	3112	1.906
53	1668	35		94	3206	1.922
54	1702	34		94	3301	1.939
55	1736	34		94	3395	1.956
56	1769	33	34.86	94	3489	1.972
57	1802	33		94	3582	1.988
58	1834	32		93	3676	2.004
59	1866	31		93	3769	2.020
60	1896	31		92	3861	2.036
61	1927	30		92	3953	2.052
62	1956	30		91	4044	2.067
63	1986	29	30.89	91	4135	2.082
64	2014	29		90	4225	2.098
65	2042	28		89	4314	2.113
66	2069	27		89	4403	2.128
67	2096	27		88	4491	2.143
68	2122	26		87	4578	2.157
69	2148	26		86	4665	2.172
70	2173	25	26.74	86	4750	2.186

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.

NOTE: In the table values are rounded, this may result in small inaccuracies when using the objectives to calculate other performance statistics.



Notes

A series of horizontal dotted lines for taking notes.



As-Hatched Performance

Day	Weight (lb) ¹	Daily gain (lb)	Av. daily gain / week (lb)	Daily intake (lb)	Cum. intake (lb) ²	FCR ³
0	0.064					
1	0.077	0.013			0.017	0.224
2	0.096	0.019		0.021	0.039	0.405
3	0.118	0.022		0.026	0.065	0.550
4	0.142	0.025		0.030	0.095	0.667
5	0.170	0.027		0.034	0.129	0.762
6	0.200	0.030		0.039	0.168	0.840
7	0.233	0.033	0.024	0.043	0.211	0.905
8	0.269	0.036		0.048	0.259	0.960
9	0.309	0.040		0.053	0.311	1.007
10	0.352	0.043		0.057	0.369	1.048
11	0.398	0.046		0.063	0.431	1.084
12	0.447	0.049		0.068	0.499	1.116
13	0.500	0.053		0.073	0.573	1.145
14	0.556	0.056	0.046	0.079	0.652	1.172
15	0.615	0.059		0.085	0.736	1.197
16	0.678	0.062		0.091	0.827	1.221
17	0.743	0.065		0.097	0.924	1.243
18	0.812	0.068		0.103	1.027	1.265
19	0.883	0.071		0.109	1.135	1.286
20	0.957	0.074		0.115	1.251	1.307
21	1.034	0.077	0.068	0.121	1.372	1.327
22	1.113	0.079		0.127	1.499	1.347
23	1.195	0.082		0.134	1.633	1.367
24	1.279	0.084		0.140	1.772	1.386
25	1.365	0.086		0.146	1.918	1.405
26	1.453	0.088		0.151	2.069	1.424
27	1.542	0.090		0.157	2.226	1.443
28	1.634	0.091	0.086	0.163	2.389	1.462
29	1.727	0.093		0.168	2.557	1.481
30	1.821	0.094		0.173	2.730	1.500
31	1.916	0.095		0.178	2.909	1.518
32	2.012	0.096		0.183	3.092	1.536
33	2.110	0.097		0.188	3.280	1.555
34	2.208	0.098		0.192	3.472	1.573
35	2.306	0.098	0.096	0.196	3.669	1.591
36	2.405	0.099		0.200	3.869	1.609
37	2.504	0.099		0.204	4.073	1.627
38	2.603	0.099		0.208	4.281	1.644
39	2.703	0.099		0.211	4.492	1.662
40	2.802	0.099		0.214	4.706	1.680
41	2.901	0.099		0.217	4.922	1.697
42	2.999	0.099	0.099	0.219	5.142	1.714
43	3.097	0.098		0.221	5.363	1.731
44	3.195	0.098		0.224	5.587	1.748
45	3.292	0.097		0.225	5.812	1.765
46	3.388	0.096		0.227	6.039	1.782
47	3.484	0.095		0.228	6.267	1.799
48	3.578	0.095		0.229	6.497	1.816
49	3.672	0.094	0.096	0.230	6.727	1.832
50	3.765	0.093		0.231	6.958	1.848
51	3.856	0.092		0.232	7.190	1.865
52	3.946	0.090		0.232	7.422	1.881
53	4.036	0.089		0.232	7.654	1.897
54	4.124	0.088		0.232	7.886	1.912
55	4.210	0.087		0.232	8.118	1.928
56	4.296	0.085	0.089	0.232	8.350	1.944
57	4.380	0.084		0.231	8.581	1.959
58	4.463	0.083		0.230	8.811	1.974
59	4.544	0.081		0.230	9.041	1.990
60	4.624	0.080		0.229	9.270	2.005
61	4.702	0.079		0.228	9.497	2.020
62	4.779	0.077		0.227	9.724	2.035
63	4.855	0.076	0.080	0.225	9.949	2.049
64	4.929	0.074		0.224	10.173	2.064
65	5.002	0.073		0.222	10.395	2.078
66	5.073	0.071		0.221	10.616	2.093
67	5.143	0.070		0.219	10.835	2.107
68	5.211	0.068		0.217	11.053	2.121
69	5.278	0.067		0.216	11.268	2.135
70	5.344	0.065	0.070	0.214	11.482	2.149

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.

NOTE: In the table values are rounded, this may result in small inaccuracies when using the objectives to calculate other performance statistics.



Rambler Ranger Broiler Performance Objectives



Male Performance

Day	Weight (lb) ¹	Daily gain (lb)	Av. daily gain / week (lb)	Daily intake (lb)	Cum. intake (lb) ²	FCR ³
0	0.064					
1	0.077	0.013			0.016	0.205
2	0.095	0.018		0.020	0.036	0.375
3	0.117	0.022		0.025	0.060	0.516
4	0.141	0.024		0.029	0.089	0.633
5	0.169	0.027		0.034	0.123	0.731
6	0.199	0.030		0.039	0.162	0.814
7	0.233	0.034	0.024	0.044	0.205	0.883
8	0.269	0.037		0.049	0.254	0.943
9	0.310	0.040		0.054	0.308	0.994
10	0.354	0.044		0.059	0.367	1.039
11	0.401	0.047		0.065	0.432	1.078
12	0.452	0.051		0.071	0.503	1.113
13	0.506	0.055		0.077	0.580	1.145
14	0.565	0.058	0.047	0.083	0.663	1.174
15	0.626	0.062		0.089	0.752	1.200
16	0.691	0.065		0.095	0.847	1.225
17	0.760	0.069		0.102	0.949	1.248
18	0.832	0.072		0.108	1.058	1.271
19	0.908	0.075		0.115	1.173	1.292
20	0.986	0.079		0.122	1.294	1.312
21	1.068	0.082	0.072	0.128	1.423	1.332
22	1.153	0.085		0.135	1.558	1.351
23	1.240	0.088		0.142	1.699	1.370
24	1.330	0.090		0.148	1.848	1.389
25	1.423	0.093		0.155	2.002	1.407
26	1.518	0.095		0.161	2.164	1.425
27	1.615	0.097		0.168	2.331	1.443
28	1.715	0.099	0.092	0.174	2.505	1.461
29	1.816	0.101		0.180	2.685	1.478
30	1.919	0.103		0.186	2.870	1.496
31	2.023	0.104		0.191	3.061	1.513
32	2.129	0.106		0.197	3.258	1.530
33	2.236	0.107		0.202	3.460	1.547
34	2.344	0.108		0.207	3.667	1.564
35	2.453	0.109	0.105	0.212	3.878	1.581
36	2.562	0.109		0.216	4.094	1.598
37	2.672	0.110		0.221	4.315	1.615
38	2.782	0.110		0.225	4.540	1.631
39	2.893	0.111		0.228	4.768	1.648
40	3.004	0.111		0.232	5.000	1.665
41	3.114	0.111		0.235	5.235	1.681
42	3.225	0.110	0.110	0.238	5.473	1.697
43	3.335	0.110		0.241	5.714	1.713
44	3.445	0.110		0.243	5.958	1.730
45	3.554	0.109		0.246	6.203	1.746
46	3.662	0.108		0.248	6.451	1.762
47	3.770	0.108		0.249	6.701	1.777
48	3.877	0.107		0.251	6.952	1.793
49	3.983	0.106	0.108	0.252	7.204	1.809
50	4.088	0.105		0.253	7.457	1.824
51	4.192	0.104		0.254	7.711	1.840
52	4.294	0.103		0.255	7.966	1.855
53	4.396	0.101		0.255	8.221	1.870
54	4.496	0.100		0.255	8.477	1.885
55	4.595	0.099		0.255	8.732	1.901
56	4.692	0.097	0.101	0.255	8.988	1.915
57	4.788	0.096		0.255	9.243	1.930
58	4.883	0.095		0.255	9.497	1.945
59	4.976	0.093		0.254	9.751	1.960
60	5.067	0.092		0.253	10.004	1.974
61	5.157	0.090		0.252	10.256	1.989
62	5.246	0.088		0.251	10.507	2.003
63	5.333	0.087	0.092	0.250	10.757	2.017
64	5.418	0.085		0.248	11.005	2.031
65	5.502	0.084		0.247	11.252	2.045
66	5.584	0.082		0.245	11.498	2.059
67	5.664	0.080		0.244	11.741	2.073
68	5.743	0.079		0.242	11.983	2.087
69	5.820	0.077		0.240	12.223	2.100
70	5.896	0.076	0.080	0.238	12.462	2.114

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.

NOTE: In the table values are rounded, this may result in small inaccuracies when using the objectives to calculate other performance statistics.



Female Performance

Day	Weight (lb) ¹	Daily gain (lb)	Av. daily gain / week (lb)	Daily intake (lb)	Cum. intake (lb) ²	FCR ³
0	0.064					
1	0.078	0.014			0.019	0.248
2	0.096	0.019		0.023	0.042	0.439
3	0.118	0.022		0.027	0.070	0.587
4	0.143	0.025		0.031	0.100	0.703
5	0.170	0.027		0.035	0.135	0.795
6	0.200	0.030		0.038	0.174	0.868
7	0.233	0.033	0.024	0.042	0.216	0.928
8	0.269	0.036		0.047	0.263	0.977
9	0.308	0.039		0.051	0.313	1.019
10	0.349	0.042		0.055	0.369	1.056
11	0.394	0.045		0.060	0.429	1.088
12	0.442	0.048		0.065	0.494	1.117
13	0.493	0.051		0.070	0.564	1.144
14	0.547	0.054	0.045	0.075	0.639	1.169
15	0.603	0.057		0.081	0.720	1.193
16	0.663	0.059		0.086	0.806	1.216
17	0.725	0.062		0.092	0.897	1.238
18	0.789	0.065		0.097	0.995	1.260
19	0.857	0.067		0.103	1.097	1.281
20	0.926	0.070		0.109	1.206	1.302
21	0.998	0.072	0.064	0.114	1.320	1.323
22	1.072	0.074		0.120	1.440	1.343
23	1.148	0.076		0.125	1.565	1.364
24	1.226	0.078		0.131	1.696	1.384
25	1.305	0.079		0.136	1.833	1.404
26	1.386	0.081		0.141	1.974	1.424
27	1.468	0.082		0.147	2.121	1.444
28	1.552	0.084	0.079	0.151	2.272	1.464
29	1.636	0.085		0.156	2.428	1.484
30	1.722	0.086		0.161	2.589	1.504
31	1.808	0.086		0.165	2.754	1.523
32	1.896	0.087		0.169	2.924	1.543
33	1.983	0.088		0.173	3.097	1.562
34	2.071	0.088		0.177	3.275	1.581
35	2.159	0.088	0.087	0.181	3.455	1.600
36	2.248	0.088		0.184	3.640	1.619
37	2.336	0.088		0.187	3.827	1.638
38	2.425	0.088		0.190	4.017	1.657
39	2.513	0.088		0.193	4.210	1.675
40	2.600	0.088		0.195	4.405	1.694
41	2.688	0.087		0.197	4.602	1.712
42	2.775	0.087	0.088	0.199	4.802	1.731
43	2.861	0.086		0.201	5.003	1.749
44	2.947	0.086		0.203	5.206	1.767
45	3.032	0.085		0.204	5.410	1.784
46	3.116	0.084		0.205	5.615	1.802
47	3.199	0.083		0.206	5.822	1.820
48	3.281	0.082		0.207	6.029	1.837
49	3.363	0.081	0.084	0.208	6.236	1.855
50	3.443	0.080		0.208	6.444	1.872
51	3.522	0.079		0.208	6.652	1.889
52	3.600	0.078		0.208	6.861	1.906
53	3.677	0.077		0.208	7.069	1.922
54	3.753	0.076		0.208	7.277	1.939
55	3.827	0.075		0.208	7.484	1.956
56	3.901	0.073	0.077	0.207	7.691	1.972
57	3.973	0.072		0.206	7.898	1.988
58	4.043	0.071		0.206	8.103	2.004
59	4.113	0.069		0.205	8.308	2.020
60	4.181	0.068		0.204	8.512	2.036
61	4.248	0.067		0.203	8.714	2.052
62	4.313	0.066		0.201	8.916	2.067
63	4.377	0.064	0.068	0.200	9.116	2.082
64	4.440	0.063		0.199	9.314	2.098
65	4.502	0.062		0.197	9.511	2.113
66	4.562	0.060		0.196	9.707	2.128
67	4.621	0.059		0.194	9.901	2.143
68	4.679	0.058		0.192	10.093	2.157
69	4.735	0.056		0.191	10.284	2.172
70	4.790	0.055	0.059	0.189	10.472	2.186

¹On-farm body weight (i.e. feed present in intestinal tract).

²Feed consumption per living bird.

³FCR includes initial body weight at placement and does not account for mortality.

NOTE: In the table values are rounded, this may result in small inaccuracies when using the objectives to calculate other performance statistics.



Notes

A series of horizontal dotted lines for taking notes.




Rambler Ranger Broiler Performance Objectives


Carcass Yield

The following diagrams indicate how yields of the major portions change with increasing live weight in each sex. Two types of processing are described: eviscerated yield is broken down into breast meat, thigh and drumstick to represent a portioning operation and into breast meat and leg meat to represent a deboning operation.

Rambler Ranger Male - Portion





Rambler Ranger Female - Portion





Rambler Ranger Broiler Performance Objectives



Definitions of Terms:

- Eviscerated %** eviscerated carcass (without neck, abdominal fat and internal organs) as a percentage of live weight.
- Breast %** breast meat (without skin and bone) as a percentage of live weight.
- Thigh / Drumstick %** whole thigh / drumstick (with skin and bone) as a percentage of live weight.
- Leg Meat %** sum of deboned thigh (without skin) and deboned drumstick (without skin) as a percentage of live weight.

NOTE: These figures represent dry yield. They do not include any moisture retained during chilling or processing. Carcass component yields will vary among processing plants depending on, for example, type of equipment used and the exact portion(s) being produced.



Notes

A series of horizontal dotted lines for taking notes.



Notes

A series of horizontal dotted lines for taking notes.



Every attempt has been made to ensure the accuracy and relevance of the information presented. However, Aviagen accepts no liability for the consequences of using the information for the management of chickens.

For further information on the management of Rowan Range® stock, please contact your local representative.

Aviagen and the Aviagen logo, the Rowan Range and the Rowan Range logo are registered trademarks of Aviagen in the US and other countries. The Rambler Ranger and the Rambler Ranger logo is a trademark of Aviagen.

All other trademarks or brands are registered by their respective owners.

© 2018 Aviagen.

0118-RR-008